

In the name of Allah, the Beneficent, the Merciful

ON THE FIELD OF DIFFERENTIAL RATIONAL INVARIANTS OF A SUBGROUP OF AFFINE GROUP(ORDINARY DIFFERENTIAL CASE)

Ural Bekbaev ¹

Department of Mathematics & Institute for Mathematical Research,
FS, UPM, 43400, Serdang, Selangor, Malaysia.

Abstract

An ordinary differential field (F, d) of characteristic zero, a subgroup H of affine group $GL(n, C) \propto C^n$ with respect to its identical representation in F^n and the following two fields of differential rational functions in $x = (x_1, x_2, \dots, x_n)$ -column vector,

$$C\langle x, d \rangle^H = \{f^d\langle x \rangle \in C\langle x, d \rangle : f^d\langle hx + h_0 \rangle = f^d\langle x \rangle \text{ for any } (h, h_0) \in H\},$$

$$C\langle x, d \rangle^{(F^*, H)} = \{f^d\langle x \rangle \in C\langle x, d \rangle : f^{g^{-1}d}\langle hx + h_0 \rangle = f^d\langle x \rangle \text{ for any } g \in F^* \text{ and } (h, h_0) \in H\}$$

are considered, where C is the constant field of (F, d) and $C\langle x, d \rangle$ is the field of differential rational functions in $x_1, x_2, \dots, x_n$ over C . The field $C\langle x, d \rangle^H$ ($C\langle x, d \rangle^{(F^*, H)}$) is an important tool in the equivalence problem of paths (respect. curves) in Differential Geometry with respect to the motion group H . In this paper an pure algebraic approach is offered to describe these fields. The field $C\langle x, d \rangle^{(F^*, H)}$ and its relation with $C\langle x, d \rangle^H$ are investigated. It is shown also that $C\langle x, d \rangle^H$ can be derived from some algebraic (without derivatives) invariants of H .

Key words: Differential field, differential rational function, invariant, differential transcendent degree.

2000 Mathematics Subject Classification: 12H05, 53A04, 53A55

1. Introduction

Let $n > 1$ be a natural number, H be a subgroup of the affine group $GL(n, R) \propto R^n$, and $G = Diff(B)$ be the group of diffeomorphisms of the open interval $B = (0, 1)$. Two infinitely smooth paths $u : B \rightarrow R^n$, $v : B \rightarrow R^n$ are said to be (G, H) -equivalent if there are such $t \in G$ and $(h, h_0) \in H$ that $v(s) = hu(t(s)) + h_0$ for any $s \in B$.

A function $f^d(u(t))$ of $u(t) = (u_1(t), \dots, u_n(t))$ and its finite number of derivatives relative to $d = \frac{d}{dt}$ is said to be invariant (more exactly, (G, H) - invariant) if the equality

$$f^d(u(t(s))) = f^\delta(hu(t(s)) + h_0)$$

is valid for any $t \in G$ and $(h, h_0) \in H$, where $\delta = \frac{d}{ds}$.

In the terms of inverse function $s = s(t)$ the expressions $\frac{du(t(s))}{dt(s)}$ and $\frac{du(t(s))}{ds}$ can be rewritten as $\frac{du(t)}{dt}$ and $\frac{du(t)}{ds(t)} = \frac{1}{s'(t)} \frac{du(t)}{dt}$, respectively. Therefore the above invariantness of $f^d(u(t))$ can be written in the form $f^d(u(t)) = f^\delta(hu(t) + h_0)$, where $\delta = s'(t)^{-1} \frac{d}{dt}$.

Let t run B , $F = C^\infty(B)$, and (F, d) be the differential ring of infinitely smooth functions on B relative to differential operator $\frac{d}{dt}$. The constant subring, i.e. $\{a \in F : d(a) = 0\}$, of (F, d) is R . Now every infinitely smooth path $u : B \rightarrow R^n$ can be considered as an element of differential module $(F^n; d)$, where d acts on elements of F^n coordinate-wisely. If elements of this module are considered as column vectors the above transformations of u and d look like

$$u = (u_1, \dots, u_n) \mapsto hu + h_0, \quad d \mapsto g^{-1}d$$

, where g is some invertible element of F and $(h, h_0) \in H$.

¹e-mail: bekbaev@science.upm.edu.my

Therefore the following algebraic approach to the above invariants is natural. Let (F, d) be any differential field i.e. F is a field with a fixed $d : F \rightarrow F$ for which

$$a) \quad d(a + b) = d(a) + d(b), \quad b) \quad d(ab) = d(a)b + ad(b) \quad \text{for any } a, b \in F.$$

Let C stand for the constant subfield of (F, d) i.e. $C = \{a \in F : d(a) = 0\}$, $H \subset GL(n, C) \propto C^n$ be a subgroup, $F^* = F \setminus \{0\}$.

One can consider the following (F^*, H) -equivalence of pairs (u, d_u) , (v, d_v) , where $u, v \in F^n$ and $d_u : F \rightarrow F$, $d_v : F \rightarrow F$ are differential operators, as an algebraic analogue of the above mentioned equivalence of paths.

Definition 1. Pairs (u, d_u) , (v, d_v) are said to be (F^*, H) -equivalent if $v = hu + h_0$, $d_v = g^{-1}d_u$ for some $g \in F^*$ and $(h, h_0) \in H$.

Let in future $x_1, \dots, x_n$ be d -differential algebraic independent variables over F and x stand for the column vector with coordinates $x_1, \dots, x_n$. We use the following notations: $C[x]$ - the ring of polynomials in variables $x_1, \dots, x_n$ over C ; $C(x)$ - the field of rational functions in variables $x_1, \dots, x_n$ over C ; $C\{x, d\}$ - the ring of differential polynomials in $x_1, \dots, x_n$ over C i.e.

$$C\{x, d\} = C[x, dx, d^2x, \dots, d^m x, \dots]$$

, where $d^m x = (d^m x_1, \dots, d^m x_n)$ and $C\langle x, d \rangle$ the field of differential rational functions in x over C .

Definition 2. An element $f^d\langle x \rangle \in C\langle x, d \rangle$ is said to be (F^*, H) invariant (F^* invariant; H invariant)- if the equality

$$f^{g^{-1}d}\langle hx + h_0 \rangle = f^d\langle x \rangle$$

(respect. $f^{g^{-1}d}\langle x \rangle = f^d\langle x \rangle$; $f^d\langle hx + h_0 \rangle = f^d\langle x \rangle$) is valid for any $g \in F^*$ and $(h, h_0) \in H$.

Let us denote the set of all (F^*, H) (respect. F^* ; H)- invariant differential rational functions over C by $C\langle x, d \rangle^{(F^*, H)}$ (respect. $C\langle x, d \rangle^{F^*}$; $C\langle x, d \rangle^H$).

Importance of differential invariants (for example such as, curvature, torsion) is out of the question. Therefore investigation of the field $C\langle x, d \rangle^{(F^*, H)}$, as far as it is an algebraic analog of such differential invariants, is of interest. In Differential Geometry usually geometric methods, for example Cartan's moving frame method, are used to obtain differential invariants of curves with respect to a motion group H . It is clear that the transformation $d \mapsto g^{-1}d$ corresponds to the ordinary change of parameter in theory of curves. In this paper we are going to offer a pure algebraic approach to describe the field $C\langle x, d \rangle^{(F^*, H)}$. Due to the algebraic character of our approach the considered case may go even beyond the current needs of Differential Geometry. The finite group case of H is considered in [1].

As to the field $C\langle x, d \rangle^H$ it is an algebraic analog of differential invariants of paths with respect to the motion group H . This field is investigated for all classical subgroups of the affine group $GL(n, R) \propto R^n$. The corresponding results and references can be found in [2].

The results of this paper are presented in [3,4].

2. Preliminary

In future $(F; d)$ stands for an ordinary differential field of characteristic zero and C is its constant field. It is assumed that the differential field (F, d) holds the following property:

If $q^d\{y\} \in F\{y, d\}$ is such a polynomial that $q^d\{a\} = 0$ for any $a \in F$ then $q^d\{y\}$ is a zero polynomial.

This condition is equivalent to $F \neq C$ due to [5, p.139].

Proposition 1. Every non constant $f^d\langle x \rangle \in F\langle x, d \rangle$ is d -algebraic independent over F .

Proof. Represent non constant $f^d\langle x \rangle$ as an irreducible ratio $\frac{P\{x\}}{Q\{x\}}$, where $P\{x\}, Q\{x\} \in F\{x, d\}$.

Assume that, for example, x_1 occurs in $P\{x\}$ or $Q\{x\}$ and define d -order of $f^d\langle x \rangle$ with respect to x_1 as the maximal k that $d^k x_1$ occurs in $P\{x\}$ or $Q\{x\}$.

To show d -algebraic independence of $f^d\langle x \rangle$ over F it is enough to show that the similar order of $df^d\langle x \rangle$ is $k + 1$. Indeed the coefficient at $d^{k+1}x_1$ in the numerator of $d\frac{P\{x\}}{Q\{x\}}$ is equal to

$$\frac{\partial P\{x\}}{\partial d^k x_1} Q\{x\} - \frac{\partial Q\{x\}}{\partial d^k x_1} P\{x\}$$

which is nonzero because characteristic of F is zero and the ring $F\{x, d\}$ is a factorial ring. Therefore for the d -order of $df^d\langle x \rangle$ with respect to x_1 one has $k + 1$.

Now let us show that if H is a subgroup then one can find such a nonzero $p^d\langle x \rangle \in C\langle x, d \rangle$ for which

$$p^{g^{-1}d}\langle hx + h_0 \rangle = g^{-1}p^d\langle x \rangle, \quad (1)$$

for any $g \in F^*$, $(h, h_0) \in H$.

Here are two examples of nonzero $p^d\langle x \rangle$ for which property (1) holds in the case of $H = GL(n, C) \propto C^n$ and therefore equality (1) is valid when H is any subgroup of the affine group as well.

Example 1. Let $g \in F^*$ and δ stand for $g^{-1}d$. The following relation can be verified.

$$d^k = \sum_{i=1}^k \Phi_{k,i}^d\{g\}\delta^i$$

, where $\Phi_{k,i}^d\{g\} = \sum C_\alpha g^{[\alpha]}$, $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k)$, $g^{[\alpha]} = g^{\alpha_1}(dg)^{\alpha_2} \dots (d^{k-1}g)^{\alpha_k}$, $C_\alpha = \frac{k!}{\alpha_1! \alpha_2! \dots \alpha_k! \prod_{j=1}^k (j!)^{\alpha_j}}$,

the sum $\sum$ is taken over all α with nonnegative integer components for which $|\alpha| = \sum_{j=1}^k \alpha_j = i$ and $1\alpha_1 + 2\alpha_2 + \dots + k\alpha_k = k$.

In future let us use the notations: $W^d = W^d\{dx\} = W^d\{dx_1, dx_2, \dots, dx_n\} = \det[dx, d^2x, \dots, d^nx]$. Let $W_i^d = W_i^d\{dx, d^{n+1}x\}$, $i = \overline{1, n}$ stand for the determinant of the matrix obtained from $[dx, d^2x, \dots, d^nx, d^{n+1}x]$ by deleting $d^i x$.

Consider one more variable y . It is easy to check that

$$W^\delta(\delta x_1, \delta x_2, \dots, \delta x_n, \delta y) = g^{-\frac{(n+1)(n+2)}{2}} W^d(dx_1, dx_2, \dots, dx_n, dy).$$

Therefore $\sum_{i=1}^{n+1} (-1)^{n+1-i} W_i^\delta \delta^i y = g^{-\frac{(n+1)(n+2)}{2}} \sum_{i=1}^{n+1} (-1)^{n+1-i} W_i^d d^i y =$

$$g^{-\frac{(n+1)(n+2)}{2}} \sum_{i=1}^{n+1} (-1)^{n+1-i} W_i^d \sum_{j=1}^i \Phi_{i,j}^d\{g\} \delta^j y = g^{-\frac{(n+1)(n+2)}{2}} \sum_{j=1}^{n+1} \left(\sum_{i=j}^{n+1} (-1)^{n+1-i} \Phi_{i,j}^d\{g\} W_i^d \right) \delta^j y.$$

It implies that for any $j = \overline{1, n+1}$ one has $W_j^\delta = g^{-\frac{(n+1)(n+2)}{2}} \sum_{i=j}^{n+1} (-1)^{i-j} \Phi_{i,j}^d\{g\} W_i^d$.

In particular, if $j = n+1$ then $W^\delta = g^{-\frac{n(n+1)}{2}} W^d$,

if $j = n$ then $W_n^\delta = g^{-\frac{n(n+1)}{2}-1} (W_n^d - \frac{n(n+1)}{2} \frac{dg}{g} W^d)$,

if $j = n-1$ then $W_{n-1}^\delta =$

$$g^{-\frac{n(n+1)}{2}-2} (W_{n-1}^d - \frac{n(n-1)}{2} \frac{dg}{g} W_n^d + (\frac{(n-1)n(n+1)}{6} d(\frac{dg}{g}) + \frac{(n-1)n(n+1)(3n-2)}{24} (\frac{dg}{g})^2) W^d),$$

if $j = n-2$ then

$$W_{n-2}^\delta = g^{-\frac{n(n+1)}{2}-3} (W_{n-2}^d - \frac{(n-1)(n-2)}{2} \frac{dg}{g} W_{n-1}^d + (\frac{(n-2)(n-1)n}{6} d(\frac{dg}{g}) +$$

$$\begin{aligned} & \frac{(n-2)(n-1)n(3n-5)}{24} \left(\frac{dg}{g}\right)^2 W_n^d - \left(\frac{(n+1)!}{24(n-3)!} d^2 \left(\frac{dg}{g}\right) + \right. \\ & \left. (2n-3) \frac{(n+1)!}{24(n-3)!} \frac{dg}{g} d\left(\frac{dg}{g}\right) + (n-1)(n-2) \frac{(n+1)!}{48(n-3)!} \left(\frac{dg}{g}\right)^3 W^d \right). \end{aligned}$$

Therefore

$$\frac{W_n^\delta}{W^\delta} = g^{-1} \left(\frac{W_n^d}{W^d} - \frac{n(n+1)}{2} \frac{dg}{g} \right) \quad (2)$$

$$\frac{W_{n-1}^\delta}{W^\delta} = g^{-2} \left(\frac{W_{n-1}^d}{W^d} - \frac{n(n-1)}{2} \frac{W_n^d}{W^d} \frac{dg}{g} + \left(\frac{(n-1)n(n+1)}{6} d\left(\frac{dg}{g}\right) + \frac{(n-1)n(n+1)(3n-2)}{24} \left(\frac{dg}{g}\right)^2 \right) \right) \quad (3)$$

Due to (2) one has the following two equalities

$$\begin{aligned} & \left(\frac{W_n^\delta}{W^\delta}\right)^2 = g^{-2} \left(\left(\frac{W_n^d}{W^d}\right)^2 - n(n+1) \left(\frac{W_n^d}{W^d} \frac{dg}{g} - \frac{n(n+1)}{4} \left(\frac{dg}{g}\right)^2 \right) \right) \\ & \delta\left(\frac{W_n^\delta}{W^\delta}\right) = g^{-2} \left(d\left(\frac{W_n^d}{W^d}\right) - \frac{W_n^d}{W^d} \frac{dg}{g} - \frac{n(n+1)}{2} \frac{d^2 g}{g} + n(n+1) \left(\frac{dg}{g}\right)^2 \right) \end{aligned} \quad (4)$$

The last equality and (3) imply that

$$\frac{W_{n-1}^\delta}{W^\delta} + \frac{n-1}{3} \delta\left(\frac{W_n^\delta}{W^\delta}\right) = g^{-2} \left(\frac{W_{n-1}^d}{W^d} + \frac{n-1}{3} d\left(\frac{W_n^d}{W^d}\right) - \frac{(n-1)(3n+2)}{6} \left(\frac{W_n^d}{W^d} \frac{dg}{g} - \frac{n(n+1)}{4} \left(\frac{dg}{g}\right)^2 \right) \right).$$

Now one can use this equality with (4) to get

$$\frac{W_{n-1}^\delta}{W^\delta} + \frac{n-1}{3} \delta\left(\frac{W_n^\delta}{W^\delta}\right) - \frac{(n-1)(3n+2)}{6n(n+1)} \left(\frac{W_n^\delta}{W^\delta}\right)^2 = g^{-2} \left(\frac{W_{n-1}^d}{W^d} + \frac{n-1}{3} d\left(\frac{W_n^d}{W^d}\right) - \frac{(n-1)(3n+2)}{6n(n+1)} \left(\frac{W_n^d}{W^d}\right)^2 \right).$$

So for the $p_1^d\langle x \rangle = \frac{W_{n-1}^d}{W^d} + \frac{n-1}{3} d\left(\frac{W_n^d}{W^d}\right) - \frac{(n-1)(3n+2)}{6n(n+1)} \left(\frac{W_n^d}{W^d}\right)^2$ one has $p_1^\delta\langle x \rangle = g^{-2} p_1^d\langle x \rangle$. Therefore $\frac{\delta p_1^\delta\langle x \rangle}{p_1^\delta\langle x \rangle} = g^{-1} \left(\frac{dp_1^d\langle x \rangle}{p_1^d\langle x \rangle} - 2 \frac{dg}{g} \right)$. Combining it with (2) one can see that the following nonzero function $p^d\langle x \rangle = \frac{n(n+1)}{2} \frac{dp_1^d\langle x \rangle}{p_1^d\langle x \rangle} - 2 \frac{W_n^d}{W^d}$ meets the needed requirements.

Example 2. Note that if one substitutes $\frac{2}{n(n+1)} \frac{W_n^d}{W^d}$ for $\frac{dg}{g}$ in the expression

$$\frac{W_{n-1}^d}{W^d} - \frac{n(n-1)}{2} \frac{W_n^d}{W^d} \frac{dg}{g} + \left(\frac{(n-1)n(n+1)}{6} d\left(\frac{dg}{g}\right) + \frac{(n-1)n(n+1)(3n-2)}{24} \left(\frac{dg}{g}\right)^2 \right),$$

which is a part of equality (3), he gets the above function $p_1^d\langle x \rangle$.

In similar way if one substitutes $\frac{2}{n(n+1)} \frac{W_n^d}{W^d}$ for $\frac{dg}{g}$ in the corresponding part of the expression for $\frac{W_{n-2}^\delta}{W^\delta}$, namely

$$\begin{aligned} & \frac{W_{n-2}^d}{W^d} - \frac{(n-1)(n-2)}{2} \frac{W_{n-1}^d}{W^d} \frac{dg}{g} + \left(\frac{(n-2)(n-1)n}{6} d\left(\frac{dg}{g}\right) + \frac{(n-2)(n-1)n(3n-5)}{24} \left(\frac{dg}{g}\right)^2 \right) \frac{W_n^d}{W^d} - \\ & \left(\frac{(n+1)!}{24(n-3)!} d^2 \left(\frac{dg}{g}\right) + (2n-3) \frac{(n+1)!}{24(n-3)!} \frac{dg}{g} d\left(\frac{dg}{g}\right) + (n-1)(n-2) \frac{(n+1)!}{48(n-3)!} \left(\frac{dg}{g}\right)^3 \right), \end{aligned}$$

then he gets a function $p_2^d\langle x \rangle$, for which $p_2^{g^{-1}d}\langle hx + h_0 \rangle = g^{-3} p_2^d\langle x \rangle$ for any $g \in F^*$ and $(h, h_0) \in H$. Therefore the function $p^d\langle x \rangle = \frac{p_2^d\langle x \rangle}{p_1^d\langle x \rangle}$ also holds the needed property (1).

To have some other examples of such $p^d\langle x \rangle$ one can see [6].

3. On (F^*, H) - invariants

Due to the above examples for a given subgroup H of the affine group $GL(n, C) \propto C^n$ one can consider $\delta(x, d) = \delta = p^d \langle x \rangle^{-1} d$, where $p^d \langle x \rangle \in C \langle x, d \rangle$ is a fixed nonzero element holding property (1). It is evident that the field $C \langle x, d \rangle^{(F^*, H)}$ is invariant with respect to the differential operator δ , so one can consider the differential field $(C \langle x, d \rangle^{(F^*, H)}, \delta)$.

Note that the field $C \langle x, d \rangle^{(F^*, H)}$ is not invariant with respect to d .

The next result deals with generators of the differential field $(C \langle x, d \rangle^{(F^*, H)}, \delta)$ over C and their relations. Let $p^d \langle x \rangle = \frac{a^d \{x\}}{b^d \{x\}}$, where $a^d \{x\}, b^d \{x\} \in C \{x, d\}$. Due to (1) for any $g \in F^*$ one has $ga^d \{x\} b^{gd} \{x\} - a^{gd} \{x\} b^d \{x\} = 0$. Consider the left side of it as a d -differential polynomial in x . In this case the above equality means that all coefficients of this polynomial, which are d -differential polynomials in g over C , are zero for any $g \in F^*$. Therefore due to assumption on (F, d) for any differential indeterminate t one has $ta^d \{x\} b^{td} \{x\} - a^{td} \{x\} b^d \{x\} = 0$. Substitution $t = p^d \langle x \rangle^{-1}$ implies that $p^\delta \langle x \rangle = 1$.

It should be noted that the field $C \langle x, d \rangle^H$ is invariant with respect to the differential operator d and $(C \langle x, d \rangle^H, d)$ is a finitely generated d -differential field over C as a subfield of $C \langle x, d \rangle$. Let us assume that $(C \langle x, d \rangle^H, d) = C \langle \varphi_1^d \langle x \rangle, \varphi_2^d \langle x \rangle, \dots, \varphi_m^d \langle x \rangle, d \rangle$. As far as $p^d \langle x \rangle \in C \langle x, d \rangle^H$ there is a differential rational function $\bar{p}^d \langle t_1, t_2, \dots, t_m \rangle$ such that

$$p^d \langle x \rangle = \bar{p}^d \langle \varphi_1^d \langle x \rangle, \varphi_2^d \langle x \rangle, \dots, \varphi_m^d \langle x \rangle \rangle.$$

Therefore due to $p^\delta \langle x \rangle = 1$ one has $\bar{p}^\delta \langle \varphi_1^\delta \langle x \rangle, \varphi_2^\delta \langle x \rangle, \dots, \varphi_m^\delta \langle x \rangle \rangle = 1$.

Theorem 1. *If $(C \langle x, d \rangle^H, d) = C \langle \varphi_1^d \langle x \rangle, \varphi_2^d \langle x \rangle, \dots, \varphi_m^d \langle x \rangle, d \rangle$ then the δ - differential field $(C \langle x, d \rangle^{(F^*, H)}, \delta)$ is generated over C by the system of elements $\varphi_1^\delta \langle x \rangle, \varphi_2^\delta \langle x \rangle, \dots, \varphi_m^\delta \langle x \rangle$. Moreover any δ -differential polynomial relation over C of the system $\varphi_1^\delta \langle x \rangle, \varphi_2^\delta \langle x \rangle, \dots, \varphi_m^\delta \langle x \rangle$ is a consequence of d -differential polynomial relations of the system $\varphi_1^d \langle x \rangle, \varphi_2^d \langle x \rangle, \dots, \varphi_m^d \langle x \rangle$ over C and the relation $\bar{p}^\delta \langle \varphi_1^\delta \langle x \rangle, \varphi_2^\delta \langle x \rangle, \dots, \varphi_m^\delta \langle x \rangle \rangle = 1$.*

Proof. It is evident that $C \langle x, d \rangle^{(F^*, H)} = C \langle x, d \rangle^{F^*} \cap C \langle x, d \rangle^H$.

Let $f^d \langle \varphi_1^d \langle x \rangle, \varphi_2^d \langle x \rangle, \dots, \varphi_m^d \langle x \rangle \rangle$ be F^* -invariant. Consider $f^{t^{-1}d} \langle \varphi_1^{t^{-1}d} \langle x \rangle, \varphi_2^{t^{-1}d} \langle x \rangle, \dots, \varphi_m^{t^{-1}d} \langle x \rangle \rangle$ as a differential rational function in x over $C \langle t, d \rangle$ and let $\frac{P_t^d \{x\}}{Q_t^d \{x\}}$ be its irreducible representation. F^* -invariantness of $f^d \langle \varphi_1^d \langle x \rangle, \varphi_2^d \langle x \rangle, \dots, \varphi_m^d \langle x \rangle \rangle$ implies that $\frac{P_t^d \{x\}}{Q_t^d \{x\}} = \frac{P_1^d \{x\}}{Q_1^d \{x\}}$. In its turn it implies that

$$P_1^d \{x\} = P_t^d \{x\} \chi^d \langle t \rangle$$

, where $\chi^d \langle t \rangle \in C \langle t, d \rangle$. Due to Proposition 1 one has $\chi^d \langle p^d \langle x \rangle \rangle \neq 0$ therefore the above equality implies that

$$\frac{P_{p^d \langle x \rangle}^d \{x\}}{Q_{p^d \langle x \rangle}^d \{x\}} = \frac{P_1^d \{x\}}{Q_1^d \{x\}} \quad \text{i.e.} \quad f^\delta \langle \varphi_1^\delta \langle x \rangle, \varphi_2^\delta \langle x \rangle, \dots, \varphi_m^\delta \langle x \rangle \rangle = f^d \langle \varphi_1^d \langle x \rangle, \varphi_2^d \langle x \rangle, \dots, \varphi_m^d \langle x \rangle \rangle$$

So it is shown that $C \langle x, d \rangle^{(F^*, H)} = C \langle \varphi_1^\delta \langle x \rangle, \varphi_2^\delta \langle x \rangle, \dots, \varphi_m^\delta \langle x \rangle, \delta \rangle$.

Now let $\psi^\delta \{ \varphi_1^\delta \langle x \rangle, \varphi_2^\delta \langle x \rangle, \dots, \varphi_m^\delta \langle x \rangle \} = 0$, where $\psi^d \{ t_1, t_2, \dots, t_m \} \in C \{ t_1, t_2, \dots, t_m, d \}$.

If $\psi^d \{ \varphi_1^d \langle x \rangle, \varphi_2^d \langle x \rangle, \dots, \varphi_m^d \langle x \rangle \} = 0$ then it means that the above relation $(\psi^\delta \{ t_1, t_2, \dots, t_m \})$ of the system $\varphi_1^\delta \langle x \rangle, \varphi_2^\delta \langle x \rangle, \dots, \varphi_m^\delta \langle x \rangle$ is a consequence of the relation $(\psi^d \{ t_1, t_2, \dots, t_m \})$ of the system $\varphi_1^d \langle x \rangle, \varphi_2^d \langle x \rangle, \dots, \varphi_m^d \langle x \rangle$ i.e. it is obtained by substitution δ for d in $\psi^d \{ t_1, t_2, \dots, t_m \}$.

If $\psi^d \{ \varphi_1^d \langle x \rangle, \varphi_2^d \langle x \rangle, \dots, \varphi_m^d \langle x \rangle \} \neq 0$ then consider $\psi^{t^{-1}d} \{ \varphi_1^{t^{-1}d} \langle x \rangle, \varphi_2^{t^{-1}d} \langle x \rangle, \dots, \varphi_m^{t^{-1}d} \langle x \rangle \}$ as a d -differential rational function in one variable t over $C \langle x, d \rangle$. Let $\frac{a_x^d \{t\}}{b_x^d \{t\}}$ be its irreducible representation and the leading coefficient (with respect to some linear order) of $b_x^d \{t\}$ be one. We show that in this case all coefficients of $a_x^d \{t\}, b_x^d \{t\}$ belong to $C \langle x, d \rangle^H$.

Indeed, first of all $\psi^{t^{-1}d}\{\varphi_1^{t^{-1}d}\langle x \rangle, \varphi_2^{t^{-1}d}\langle x \rangle, \dots, \varphi_m^{t^{-1}d}\langle x \rangle\}$, as a differential rational function in x , is H -invariant function, as much as $\varphi_i^d\langle x \rangle \in C\langle x, d \rangle^H$. This H -invariantness implies that

$$a_x^d\{t\}b_{hx+h_0}^d\{t\} = b_x^d\{t\}a_{hx+h_0}^d\{t\}$$

for any $(h, h_0) \in H$. Therefore $b_{hx+h_0}^d\{t\} = \chi^d\langle x, (h, h_0) \rangle b_x^d\{t\}$. But comparison of the leading terms of both sides implies that in reality $\chi^d\langle x, (h, h_0) \rangle = 1$ which in its turn implies that all coefficients of $b_x^d\{t\}$ (as well as $a_x^d\{t\}$) are H -invariant.

Therefore all coefficients of $a_x^d\{t\}$, $b_x^d\{t\}$ can be considered as d -differential rational functions in $\varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle$, for example $b_x^d\{t\} = \bar{b}_{\varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle}\{t\}$. Now represent the numerator $a_x^d\{t\}$ as a d -differential polynomial function in $t - \bar{p}^d\langle \varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle \rangle$, for example $a_x^d\{t\} = \bar{a}_{\varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle}^d\{t - \bar{p}^d\langle \varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle \rangle\}$. As such polynomial its constant term is zero because of $\psi^\delta\{\varphi_1^\delta\langle x \rangle, \varphi_2^\delta\langle x \rangle, \dots, \varphi_m^\delta\langle x \rangle\} = 0$. So

$$\psi^{t^{-1}d}\{\varphi_1^{t^{-1}d}\langle x \rangle, \varphi_2^{t^{-1}d}\langle x \rangle, \dots, \varphi_m^{t^{-1}d}\langle x \rangle\} = \frac{\bar{a}_{\varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle}^d\{t - \bar{p}^d\langle \varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle \rangle\}}{\bar{b}_{\varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle}^d\{t\}}.$$

Substitution $t = 1$ implies that

$$\psi^d\{\varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle\} = \frac{\bar{a}_{\varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle}^d\{1 - \bar{p}^d\langle \varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle \rangle\}}{\bar{b}_{\varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle}^d\{1\}}.$$

Now consider the following δ -differential rational function over C :

$$\bar{\psi}^\delta\langle t_1, t_2, \dots, t_m \rangle = \psi^\delta\{t_1, t_2, \dots, t_m\} - \frac{\bar{a}_{t_1, t_2, \dots, t_m}^\delta\{1 - \bar{p}^\delta\langle t_1, t_2, \dots, t_m \rangle\}}{b_{t_1, t_2, \dots, t_m}^\delta\{1\}}.$$

For this function one has

$$\bar{\psi}^\delta\langle \varphi_1^\delta\langle x \rangle, \varphi_2^\delta\langle x \rangle, \dots, \varphi_m^\delta\langle x \rangle \rangle = 0 \quad \text{as well as} \quad \bar{\psi}^d\langle \varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle \rangle = 0.$$

But once again the last equality (relation) means that it is consequence of relations of the system $\varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle, \dots, \varphi_m^d\langle x \rangle$. This is the end of proof of Theorem 1.

Example 3. Let $C = R$, $n = 2$ and

$$H = O(2, R) = \{h \in GL(2, R) : hh^t = E\}$$

In this case $R\langle x, d \rangle^H = R\langle \varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle; d \rangle$, where $\varphi_1^d\langle x \rangle = (x, x)$, $\varphi_2^d\langle x \rangle = (dx, dx)$ and $(*, *)$ stands for the dot product. Moreover $\varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle$ is d -algebraic independent over R . For $p^d\langle x \rangle$ one can take (x, dx) , so $\delta = (x, dx)^{-1}d$. Due to the Theorem 1

$$(R\langle x, d \rangle^{(F^*, H)}, \delta) = R\langle \varphi_1^\delta\langle x \rangle, \varphi_2^\delta\langle x \rangle; \delta \rangle$$

In this case $\bar{p}^d\langle \varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle \rangle = \frac{1}{2}d\varphi_1^d\langle x \rangle$. So $p^\delta\langle x \rangle = \frac{1}{2}\delta\varphi_1^\delta\langle x \rangle = \frac{1}{2}\frac{1}{(x, dx)}d(x, x) = 1$ and $\varphi_2^\delta\langle x \rangle = (\delta x, \delta x)$ is δ -algebraic independent over R .

Example 4. Let now $H = O(2, R) \propto R^2$. In this case $R\langle x, d \rangle^H = R\langle \varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle; d \rangle$, where $\varphi_1^d\langle x \rangle = (dx, dx)$, $\varphi_2^d\langle x \rangle = (d^2x, d^2x)$, and $\varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle$ is d -algebraic independent over R . For $p^d\langle x \rangle$ one can take

$$d \frac{\det[dx, d^2x]^2}{(dx, dx)^3} = d \frac{\varphi_1^d\langle x \rangle \varphi_2^d\langle x \rangle - \frac{1}{4}(d\varphi_1^d\langle x \rangle)^2}{(\varphi_1^d\langle x \rangle)^3} = \bar{p}^d\langle \varphi_1^d\langle x \rangle, \varphi_2^d\langle x \rangle \rangle$$

Due to the Theorem 1

$$(R\langle x, d \rangle^{(F^*, H)}, \delta) = R\langle \varphi_1^\delta \langle x \rangle, \varphi_2^\delta \langle x \rangle; \delta \rangle$$

In this case we have one basic relation

$$\bar{p}^\delta \langle \varphi_1^\delta \langle x \rangle, \varphi_2^\delta \langle x \rangle \rangle = 1$$

The next important problem is the finding out the number of elements of a maximal δ -algebraic independent system of elements of $C\langle x, d \rangle^{(F^*, H)}$ over C i.e. evaluation the differential transcendence degree $\delta - \text{tr.deg} C\langle x, d \rangle^{(F^*, H)} / C$ of $C\langle x, d \rangle^{(F^*, H)}$ over the its field of constants C . Theorem 2 deals with this problem.

Theorem 2. $\delta - \text{tr.deg} C\langle x, d \rangle^{(F^*, H)} / C = n - 1$.

Proof. Indeed due to Theorem 1, applied to H consisting of only identity element, $C\langle x, d \rangle^{F^*} = C\langle x, \delta \rangle$ with only one main relation among $x_1, x_2, \dots, x_n$, namely $p^\delta \langle x \rangle = 1$ i. e. $a^\delta \{x\} - b^\delta \{x\} = 0$. But $a^\delta \{x\} - b^\delta \{x\}$ is not zero polynomial as far as $p^\delta \langle x \rangle \neq 1$. Therefore $\delta - \text{tr.deg} C\langle x, \delta \rangle / C = n - 1$.

Now to establish the equality $\delta - \text{tr.deg} C\langle x, d \rangle^{(F^*, H)} / C = n - 1$ it is enough to show that every x_i is δ -differential algebraic over $(C\langle x \rangle^{(F^*, H)}, \delta)$. For it consider the following

$$W^\delta(\delta x_1, \delta x_2, \dots, \delta x_n)^{-1} W^\delta(\delta x_1, \delta x_2, \dots, \delta x_n, \delta y) = 0$$

as a differential equation in one variable y .

One can check easily that

$$W^\delta(\delta x_1, \delta x_2, \dots, \delta x_n)^{-1} W^\delta(\delta x_1, \delta x_2, \dots, \delta x_n, \delta y) = \sum_{i=1}^{n+1} (-1)^{n+1-i} \frac{W_i^\delta}{W^\delta} \delta^i y$$

But $\frac{W_i^\delta}{W^\delta} \in C\langle x, d \rangle^{(F^*, H)}$ and $y = x_i$ is a solution of the above differential equation for any $i = \overline{1, n}$, which implies that $\delta - \text{tr.deg} C\langle x, d \rangle^{(F^*, H)} / C = n - 1$. This is the end of proof of Theorem 2.

4. On differential field $(C\langle x, d \rangle^H, d)$

Due to Theorem 1 one can find a system of differential generators of $(C\langle x, d \rangle^H, d)$ to get a system of differential generators of $(C\langle x, d \rangle^{(F^*, H)}, \delta)$. In [2] it is proved that $C\langle x, d \rangle^{GL(n, C) \ltimes C^n}$ as a d -differential field over C is generated by the following d -algebraic independent system $\frac{W_1^d}{W^d}, \frac{W_2^d}{W^d}, \dots, \frac{W_n^d}{W^d}$.

The next result reduces the problem of description of the differential field $(C\langle x, d \rangle^H, d)$ to the description of the ordinary field of algebraic invariants of H .

Theorem 3. *The system $x, dx, d^2x, \dots, d^n x$ is algebraic independent over $C\langle x, d \rangle^{GL(n, C) \ltimes C^n}$ and $C\langle x, d \rangle = C\langle x, d \rangle^{GL(n, C) \ltimes C^n}(x, dx, d^2x, \dots, d^n x)$.*

Proof. It is evident that it is enough to prove the algebraic independence of the system $dx, d^2x, \dots, d^n x$ over $C\langle x, d \rangle^{GL(n, C) \ltimes C^n}$. Let $P[z_1, z_2, \dots, z_n]$, where $z_i = (z_{1i}, z_{2i}, \dots, z_{ni})$, be a nonzero polynomial over $C\langle x, d \rangle^{GL(n, C) \ltimes C^n}$ such that $P[dx, d^2x, \dots, d^n x] = 0$. Assume, for example, at least one of $z_{n1}, z_{n2}, \dots, z_{nn}$ occurs in P and

$$P[z_1, z_2, \dots, z_n] = \sum_{\alpha} (z_{n1})^{\alpha_1} (z_{n2})^{\alpha_2} \dots (z_{nn})^{\alpha_n} P_{\alpha}[\bar{z}_1, \bar{z}_2, \dots, \bar{z}_n]$$

, where $P_{\alpha}[\bar{z}_1, \bar{z}_2, \dots, \bar{z}_n]$ are polynomials over $C\langle x, d \rangle^{GL(n, C) \ltimes C^n}$ in $\bar{z}_i = (z_{1i}, z_{2i}, \dots, z_{(n-1)i})$, $i = \overline{1, n}$. To prove Theorem 3 it is enough to show that $P[dx, d^2x, \dots, d^n x] = 0$ implies $P_{\alpha}[\bar{dx}, \bar{d^2x}, \dots, \bar{d^nx}] = 0$ for all α .

Consider $h \in GL(n, C)$ which's i -th column is of the form $(0, \dots, 0, 1, 0, \dots, 0, c_i)$, where $i = \overline{1, n-1}$ and its n -th column is $(0, \dots, 0, c_n)$. For such h one has $\bar{d^i}(hx) = \bar{d^i}x$. So far as the coefficients of

$P[z_1, z_2, \dots, z_n]$ are $GL(n, C) \propto C^n$ -invariant, substitution hx for x into $P[dx, d^2x, \dots, d^nx] = 0$ implies that

$$\sum_{\alpha} \left(\sum_{i=1}^n c_i dx_i \right)^{\alpha_1} \left(\sum_{i=1}^n c_i d^2x_i \right)^{\alpha_2} \dots \left(\sum_{i=1}^n c_i d^nx_i \right)^{\alpha_n} P_{\alpha}[\overline{dx}, \overline{d^2x}, \dots, \overline{d^nx}] = 0$$

Therefore due to the assumption on (F, d) for variables $y_1, y_2, \dots, y_n$ one has

$$\sum_{\alpha} \left(\sum_{i=1}^n y_i dx_i \right)^{\alpha_1} \left(\sum_{i=1}^n y_i d^2x_i \right)^{\alpha_2} \dots \left(\sum_{i=1}^n y_i d^nx_i \right)^{\alpha_n} P_{\alpha}[\overline{dx}, \overline{d^2x}, \dots, \overline{d^nx}] = 0 \quad (5)$$

Now consider the ring $C\langle x, d \rangle[y_1, y_2, \dots, y_n]$ with respect to differential operators $\partial_1 = \frac{\partial}{\partial y_1}, \partial_2 = \frac{\partial}{\partial y_2}, \dots, \partial_n = \frac{\partial}{\partial y_n}$. It is clear that its constant ring is $C\langle x, d \rangle$ i.e.

$$C\langle x, d \rangle = \{a \in C\langle x, d \rangle[y_1, y_2, \dots, y_n] : \partial_1 a = \partial_2 a = \dots \partial_n a = 0\}.$$

Introduce new differential operators $\overline{\partial}_i = \sum_{j=1}^n f_{ij}^d \langle x \rangle \partial_j$, where $i = \overline{1, n}$,

$$(f_{ij}^d \langle x \rangle)_{i,j=\overline{1,n}} = [dx, d^2x, \dots, d^nx]^{-1}$$

The following are evident:

a) The constant ring of $C\langle x, d \rangle[y_1, y_2, \dots, y_n]$ with respect to new differential operators $\overline{\partial}_1, \overline{\partial}_2, \dots, \overline{\partial}_n$ is the same $C\langle x, d \rangle$,

b) $\overline{\partial}_j(\sum_{i=1}^n y_i d^k x_i)$ is equal to 0 whenever $j \neq k$ and it is equal to 1 if $j = k$.

Now if one assumes that $\alpha^0 = (\alpha_1^0, \dots, \alpha_n^0)$ is a such one for which

$$|\alpha^0| = \max\{|\alpha| : P_{\alpha}[\overline{dx}, \overline{d^2x}, \dots, \overline{d^nx}] \neq 0\}$$

and applies $(\overline{\partial}_1)^{\alpha_1^0} (\overline{\partial}_2)^{\alpha_2^0} \dots (\overline{\partial}_n)^{\alpha_n^0}$ to equality (5) he comes to a contradiction $P_{\alpha^0}[\overline{dx}, \overline{d^2x}, \dots, \overline{d^nx}] = 0$.

The equality $C\langle x, d \rangle = C\langle x, d \rangle^{GL(n, C) \propto C^n}(x, dx, \dots, d^nx)$ is an easy consequence of the fact that $y = x_i, i = \overline{1, n}$, are solutions of the differential equation

$$W(dx_1, dx_2, \dots, dx_n)^{-1} W(dx_1, dx_2, \dots, dx_n, dy) = 0$$

all coefficients of which belong to $C\langle x, d \rangle^{GL(n, C) \propto C^n}$. This is the end of proof Theorem 3.

So due to Theorem 3

$$C\langle x, d \rangle^H = C\langle x, d \rangle^{GL(n, C) \propto C^n}(x, dx, \dots, d^nx)^H$$

and the system $\{x, dx, \dots, d^nx\}$ is algebraic independent over $C\langle x, d \rangle^{GL(n, C) \propto C^n}$. But as we have already noticed the d -differential field $C\langle x, d \rangle^{GL(n, C) \propto C^n}$ is generated over C by the d -algebraic independent system $\frac{W_1^d\{dx, d^{n+1}x\}}{W^d\{dx\}}, \frac{W_2^d\{dx, d^{n+1}x\}}{W^d\{dx\}}, \dots, \frac{W_n^d\{dx, d^{n+1}x\}}{W^d\{dx\}}$. Therefore to find a system of generators of the field $C\langle x, d \rangle^{GL(n, C) \propto C^n}(x, dx, \dots, d^nx)^H$ over $C\langle x, d \rangle^{GL(n, C) \propto C^n}$ it is enough to find a system of generators of the field $C(x, dx, \dots, d^nx)^H$ over C (see the Appendix). Due to these the following method can be used to construct a system of generators of the d -differential field $C\langle x, d \rangle^H$ over C :

1) Find any system of ordinary algebraic generators of the field $C(z_1, z_2, \dots, z_{n+1})^H$, where $z_i = (z_{1i}, z_{2i}, \dots, z_{ni})$, $i = \overline{1, n+1}$, and the action of H is defined as:

$$((h, h_0), (z_1, z_2, \dots, z_{n+1})) \rightarrow (hz_1 + h_0, hz_2, \dots, hz_{n+1})$$

For example, let it be $\varphi_1(z_1, z_2, \dots, z_{n+1}), \varphi_2(z_1, z_2, \dots, z_{n+1}), \dots, \varphi_k(z_1, z_2, \dots, z_{n+1})$.

2) The system

$$\frac{W_1^d\{dx, d^{n+1}x\}}{W^d\{dx\}}, \frac{W_2^d\{dx, d^{n+1}x\}}{W^d\{dx\}}, \dots, \frac{W_n^d\{dx, d^{n+1}x\}}{W^d\{dx\}}, \varphi_1(x, dx, \dots, d^nx), \dots, \varphi_k(x, dx, \dots, d^nx)$$

will be a system of generators of the differential field $(C\langle x, d \rangle^H, d)$ over C .

5. Discussion of two questions

The following kind questions are typical in Differential Geometry. The complete answers to these questions can not be given in our case because of its generality. The answers depend on differential field (F, d) and group H .

The first one is if $C\langle x, d \rangle^{(F^*, H)}$ distinguishes nonequivalent curves ("of common position"). In our case it should be noted that if pairs $(u, d_1), (v, d_2)$ are (F^*, H) -equivalent then $\delta(u, d_1) = \delta(v, d_2)$ i.e. $(p^{d_1}\langle u \rangle)^{-1}d_1 = (p^{d_2}\langle v \rangle)^{-1}d_2$, provided that $p^{d_1}\langle u \rangle, p^{d_2}\langle v \rangle$ are not zero. Therefore in our case the equality $\delta_0 = (p^{d_1}\langle u \rangle)^{-1}d_1 = (p^{d_2}\langle v \rangle)^{-1}d_2$ is a natural condition for the pairs $(u, d_1), (v, d_2)$ to be (F^*, H) -equivalent.

Consider $X = \{v \in F^n : W^d\{dv\}p^d\langle v \rangle \neq 0\}$, which is not empty due to the assumption on differential field (F, d) . Assume that $(u, d_1), (v, d_2)$ are such pairs that $\delta_0 = \delta(u, d_1) = \delta(v, d_2)$ and for every $\varphi^\delta\langle x \rangle \in C\langle x, d \rangle^{(F^*, H)}$ the equality $\varphi^{\delta_0}\langle u \rangle = \varphi^{\delta_0}\langle v \rangle$ is valid. Then in particular

$$a_i = \frac{W_i^{\delta_0}\{\delta_0 u, \delta_0^{n+1} u\}}{W^{\delta_0}\{\delta_0 u\}} = \frac{W_i^{\delta_0}\{\delta_0 v, \delta_0^{n+1} v\}}{W^{\delta_0}\{\delta_0 v\}}$$

at any $i = \overline{1, n}$. Which means that components both of u, v are solutions of the same $(n+1)$ -order linear differential equation

$$\delta_0^{n+1}y + \sum_{i=1}^n (-1)^{n+1-i} a_i \delta_0^i y = 0$$

one more solution of which is $y = 1$. It means that $v = hu + h_0$ for some $(h, h_0) \in GL(n, C) \propto C^n$. Therefore in common case, the problem is reduced to the question if $C(z_1, z_2, \dots, z_{n+1})^H$ distinguishes $u, du, \dots, d^n u$ and $hu + h_0, hdu, \dots, h d^n u$.

Another question is: What values can take a given system generators of $(C\langle x, d \rangle^{(F^*, H)}, \delta)$? Let us consider the case when the system of generators of $(C\langle x, d \rangle^H, d)$ is given in the above considered form:

$$\frac{W_i^d\{dx, d^{n+1}x\}}{W^d\{dx\}}, i = \overline{1, n}, \quad \varphi_j(x, dx, \dots, d^n x), j = \overline{1, k}$$

and let us assume that $\{\psi_i^d\{y_1, y_2, \dots, y_n, t_1, t_2, \dots, t_k\}_{i=\overline{1, l}}\}$ is a fundamental system of relations of the above system i.e. any other relation is an element of the radical differential ideal generated by this system in $C\{y_1, y_2, \dots, y_n, t_1, t_2, \dots, t_k; d\}$. Then due to Theorem 1 the system

$$\frac{W_i^\delta\{\delta x, \delta^{n+1}x\}}{W^\delta\{\delta x\}}, i = \overline{1, n}, \quad \varphi_j(x, \delta x, \dots, \delta^n x), j = \overline{1, k}$$

is a system of generators of the differential field $(C\langle x, d \rangle^{(F^*, H)}, \delta)$ and the system consisting of $\overline{p}^\delta\langle y_1, y_2, \dots, y_n, t_1, t_2, \dots, t_k \rangle - 1, \psi_i^\delta\{y_1, y_2, \dots, y_n, t_1, t_2, \dots, t_k\}$, where $i = \overline{1, l}$, is a fundamental system of relations of this system of generators.

Therefore if $(a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_k) \in F^{n+k}$ is a fixed element then the question if one can find such $u = (u_1, u_2, \dots, u_n) \in F^n$ for which $\frac{W_i^{\delta_0}\{\delta_0 u, \delta_0^{n+1} u\}}{W^{\delta_0}\{\delta_0 u\}} = a_i$, where $i = \overline{1, n}$ and $\varphi_j(u, \delta_0 u, \dots, \delta_0^n u) = b_j$, where $j = \overline{1, k}$, $\delta_0 = p^d\langle u \rangle^{-1}d$, is equivalent to the existence of a solution in F^n of the following system:

$$\begin{cases} \psi_i^\delta\{a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_k\} = 0, & \text{where } i = \overline{1, l} \\ \overline{p}^\delta\langle a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_k \rangle - 1 = 0 \\ \delta^{n+1}x + \sum_{i=1}^n (-1)^{n+1-i} a_i \delta^i x = 0 \\ \varphi_j(x, \delta x, \dots, \delta^n x) = b_j, & \text{where } j = \overline{1, k} \end{cases}$$

and, of course, $\delta = p^d \langle x \rangle^{-1} d$.

6. Appendix

Let C be any field of characteristic zero, t and z , where z stands for column vector with coordinates $z_1, \dots, z_n$, are indeterminates over C , H be any subgroup of $GL(n, C) \propto C^n$. For brevity, let us denote an element of H as h and its action on z as hz . We consider $C(t, z)$ with respect to transformations $(h; (t, z)) \rightarrow (t, hz)$.

Proposition 2. *The field $C(t, z)^H$ is generated by $C(z)^H$ over $C(t)$.*

Proof. Let $\frac{P[z]}{Q[z]}$ be any irreducible ratio from $C(t, z)^H$. One can assume that coefficients of $P[z], Q[z]$ are from $C[t]$. For any $h \in H$ the equality $\frac{P[z]}{Q[z]} = \frac{P[hz]}{Q[hz]}$ implies that

$$P[hz] = P[z]\chi_h[z] \quad (6)$$

, where $\chi_h[z] \in C(t)[z]$. In particular $P[h^{-1}z] = P[z]\chi_{h^{-1}}[z]$ i.e. $P[z] = P[hz]\chi_h[hz]$. Combining it with (6) implies that $P[hz] = P[hz]\chi_h[z]\chi_{h^{-1}}[hz]$ i.e. $\chi_h[z]\chi_{h^{-1}}[hz] = 1$. It shows that $\chi_h[z]$ does not depend on z , i.e. $\chi_h[z] \in C(t)$. Now consider the leader z^α of $P[z]$, with respect to any linear order, and apply $(\frac{\partial}{\partial z_1})^{\alpha_1} \dots (\frac{\partial}{\partial z_n})^{\alpha_n}$ to equality (6). It shows that in reality $\chi_h[z]$ is of the form $\chi_h[z] = \frac{\phi_h[t]}{a[t]}$, $\phi_h[t] = \sum_{i=0}^k \phi_i(h)t^i$, where k is a nonnegative integer, $\phi_i(h) \in C(h)$, $a[t] \in C[t]$, the leading coefficient of $a[t]$ is assumed to be 1. Due to (6) the polynomial $\phi_h[t]$ has property

$$\phi_{h_1}[t]\phi_{h_2}[t] = \phi_{h_1 h_2}[t]a[t] \quad (7)$$

for any $h_1, h_2 \in H$. In particular it implies that

$$\phi_k(h_1)\phi_k(h_2) = \phi_k(h_1 h_2)$$

and therefore if $\phi_k[h_1] \neq 0$ for some $h_1 \in H$ then $\phi_k[h] \neq 0$ for any $h \in H$. Now due to (7) it is clear that $\deg \phi_h[t] = \deg a[t]$ for any $h \in H$. But due to (7) if some prime $p[t] \in C[t]$ divides $a[t]$ then it divides $\phi_h[t]$ as well. It is not difficult to see now that $a[t]$ divides $\phi_h[t]$ and therefore $\phi_h[t] = \phi_k(h)a[t]$ i.e. $\chi_h[z] = \frac{\phi_h[t]}{a[t]} = \phi_k(h)$. Represent $P[z]$ ($Q[z]$) in the form $\sum P_i[z]t^i$ (respect. $\sum Q_i[z]t^i$), where $P_i[z] \in C[z]$ (respect. $Q_i[z] \in C[z]$), rewrite (6) in the following form $\sum P_i[hz]t^i = \sum P_i[z]t^i \phi_k(h)$ (respect. $\sum Q_i[hz]t^i = \sum Q_i[z]t^i \phi_k(h)$). So for every i one has $P_i[hz] = P_i[z]\phi_k(h)$ (respect. $Q_i[hz] = Q_i[z]\phi_k(h)$) and therefore $\frac{P[z]}{Q[z]} \in C(t)C(z)^H$. This is the end of proof.

References

1. U.D. Bekbaev, On differential rational invariants of finite subgroups of Affine group. Bulletin of Malaysian Mathematical Society, 2005, Volume 28, N1, pp. 55-60.
2. Dj.Khadjiev, Application of Invariant Theory to Differential Geometry of curves, FAN, Tashkent, 1988, (Russian).
3. U.D. Bekbaev, On invariants of Curves, INSPEM, Technical report No.1 December, 2002
4. U.D. Bekbaev, On invariants of curves. Proceedings of International Conference on Research and Education in Mathematics, 2-4 April 2003. At: Hotel Equatorial, Kuala Lumpur, Malaysia. pp. 82-95. University Putra Malaysia Press.
5. E.R. Kolchin, Differential Algebra and Algebraic Groups, Academic Press, New York, 1973.
6. U.D. Bekbaev, Once again on equivalence and invariants of differential equations $y^{(n)} + a_{n-1}y^{(n-1)} + \dots + a_0y = 0$, Uzbek Mathematical Journal. 3(1995)19-31 (Russian).